

How Phase-Breaking Affects Quantum Transport Through Chaotic Cavities

Harold U. Baranger¹ and Pier A. Mello²

¹ AT&T Bell Laboratories 1D-230, 600 Mountain Avenue, Murray Hill, New Jersey 07974-0636

² Instituto de Física, Universidad Nacional Autónoma de México, 01000 México D.F., México

(Submitted to Phys. Rev. B, November 14, 1994)

We investigate the effects of phase-breaking events on electronic transport through ballistic chaotic cavities. We simulate phase-breaking by a fictitious lead connecting the cavity to a phase-randomizing reservoir and introduce a statistical description for the total scattering matrix, including the additional lead. For strong phase-breaking, the average and variance of the conductance are calculated analytically. Combining these results with those in the absence of phase-breaking, we propose an interpolation formula, show that it is an excellent description of random-matrix numerical calculations, and obtain good agreement with several recent experiments.

PACS numbers: 72.20.My, 05.45.+b, 72.15.Gd

Recently there has been great interest in the effects of quantum-mechanical interference on electronic transport through ballistic quantum dots.¹ In these microstructures both the phase-coherence length and the elastic mean free path exceed the system dimensions. Thus the leads into the dot can be thought of as electron waveguides and the dot itself as a resonant cavity.

Experimentally^{2,3,4,5,6,7,8,9} one observes random (but reproducible) fluctuations in the conductance as the magnetic field,^{2,4,9} the Fermi energy,⁵ or the shape of the cavity⁹ is changed. The sensitivity to small changes in these parameters shows that these fluctuations are caused by quantum interference. In addition to conductance fluctuations, several interference effects which survive averaging over many microstructures are observed;^{6,7,8} in particular, an increase in the average resistance at zero magnetic field called the weak-localization correction (WLC).⁸ The two main features of these interference effects are their *shape*— the characteristic field or energy to which they are sensitive, or more precisely the functional dependence on these parameters— and their *magnitude*— simply how big these quantum corrections are compared to the classical conductance. Here we concentrate on the magnitude and merely note that a theory for the shape has been extensively developed.^{10,11,12,13}

A theory of the magnitude of quantum interference effects in chaotic cavities has recently been developed by making a statistical ansatz for the S -matrix describing the scattering.^{12,13,14,15,16} Refs. 15 and 16 developed a random S -matrix theory by assigning to S an “equal a priori distribution” once the symmetry requirements were imposed. The results for the average, variance, and probability density of the conductance were in good agreement with numerical calculations for a chaotic cavity connected to two waveguides.¹⁵ However, the random-matrix predictions for both the weak-localization correction and the variance are larger than the experimental results.^{8,9} In addition, the measured probability density is close to a Gaussian distribution⁹ when there are two

propagating modes per lead ($N = 2$), while random-matrix theory predicts a Gaussian distribution only for $N \geq 3$.¹⁵

Inherent in the model of Refs. 15 and 16 is the assumption (among others) that one can neglect phase-breaking processes which destroy the coherence of the wave function. In this paper we show that this assumption is largely responsible for the discrepancy between theory and experiment mentioned above. We make specific predictions for the dependence of the quantum transport corrections on the degree of phase-breaking which may be tested by future experiments.

To simulate the effects of phase-breaking events we adopt a model suggested by M. Büttiker:¹⁷ in addition to the physical leads 1, 2 attached to reservoirs at chemical potentials μ_1, μ_2 , a lead 3 connects the cavity to a phase-randomizing reservoir at μ_3 . This model has been discussed extensively for disordered materials¹⁸ and, more recently, for ballistic quantum dots by Marcus, *et al.*³ A similar model has been used for absorption of microwaves in chaotic-scattering from cavities.¹⁹ Requiring the current in lead 3 to vanish determines μ_3 ; the two-terminal dimensionless conductance is then found to be

$$g \equiv G/(e^2/h) = 2 \left[T_{21} + \frac{T_{23}T_{31}}{T_{32} + T_{31}} \right], \quad (1)$$

where T_{ij} is the transmission coefficient for “spinless electrons” from lead j to lead i . The factor of 2 accounts for spin explicitly. We call N the number of channels in leads 1 and 2, N_ϕ that in lead 3, and $N_T = 2N + N_\phi$.

We now make the fundamental assumption of an “equal a priori distribution” for the *total* $N_T \times N_T$ scattering matrix S , once the symmetry requirements have been imposed. S is, of course, unitary, and is symmetric in the absence of a magnetic field because of time-reversal symmetry. We assume that the statistics of the total S -matrix are given by the Circular Ensembles of random matrix theory.²⁰ For $B = 0$ the orthogonal ensemble (COE, denoted $\beta = 1$) is appropriate while for

nonzero B we use the unitary ensemble (CUE, $\beta = 2$). In contrast to previous studies of the eigenphases,^{10,16} total transmission,^{15,16} or individual S -matrix elements,¹⁹ we treat the statistics of g given in Eq. (1). In the rest of the paper, we first derive results valid in the weak and strong phase-breaking limits, then combine these into an interpolation formula which simulations show to be valid, and finally compare with experiments.

We start by recalling the result for the WLC and variance at $N_\phi = 0$ given in Ref. 15:

$$\delta g \equiv \langle g \rangle^{(\beta=1)} - \langle g \rangle^{(\beta=2)} = -N/(2N+1) \quad (2a)$$

$$\text{varg} = \begin{cases} \frac{4N(N+1)^2}{(2N+1)^2(2N+3)}, & \beta = 1 \\ \frac{N^2}{4N^2-1}, & \beta = 2. \end{cases} \quad (2b)$$

In addition to these results, the probability density of g was calculated for $N = 1 - 3$.

The case $N_\phi = 1$ and $\beta = 1$ is a special one which can be analyzed in detail. From the joint probability distribution of S -matrix elements in one row,²¹

$$P(S_{11}, \dots, S_{1N_T}) \propto (1 - |S_{11}|^2)^{-N} \delta(1 - \sum_{i=1}^{N_T} |S_{1i}|^2), \quad (3)$$

one can show that the joint distribution of the T_{3j} 's is

$$P(\{T_{3j}\}) = \frac{N(2N-1)!(T_{31}T_{32})^{N-1} \delta(1 - \sum T_{3j})}{[(N-1)!]^2 (T_{31} + T_{32})^N}. \quad (4)$$

Remarkably, the $\langle g \rangle$ that one obtains by integrating over this distribution is identical to that for $N_\phi = 0$.

In the limit $N_\phi \gg 1$, one can obtain a number of results to leading order in $1/N_\phi$. First, expand the conductance of Eq. (1) in powers of δT_{ij} where $T_{ij} = \langle T_{ij} \rangle + \delta T_{ij}$. By definition $\langle \delta T_{ij} \rangle = 0$; using the methods of Ref. 22, one finds that the correlations among the T_{ij} 's are given by

$$\begin{aligned} \langle \delta T_{ij} \delta T_{kl} \rangle &= A_\beta \{ N_i N_j (N_T + \delta_{\beta 1}) (N_T + 2\delta_{\beta 1}) \delta_{ik} \delta_{jl} \\ &\quad - N_i N_j N_l (N_T + \delta_{\beta 1}) \delta_{ik} - N_i N_j N_k (N_T + \delta_{\beta 1}) \delta_{jl} \\ &\quad + N_i N_j N_k N_l (1 + \delta_{\beta 1}) \}, \end{aligned} \quad (5a)$$

$$A_1 = [N_T(N_T+1)^2(N_T+3)]^{-1}, \quad (5b)$$

$$A_2 = [N_T^2(N_T^2-1)]^{-1} \quad (5c)$$

for $i \neq j, k \neq l$. (For $\beta = 1$, the indices of the T_{ij} 's have been permuted so as to maximize coincidences.) Note that $\langle \delta T_{ij} \delta T_{kl} \rangle$ is at least of order $1/N_\phi^2$. Thus

$$\langle g \rangle = 2 \left[\langle T_{21} \rangle + \frac{\langle T_{23} \rangle \langle T_{31} \rangle}{\langle T_{32} \rangle + \langle T_{31} \rangle} \right] + O(1/N_\phi^2). \quad (6)$$

Here we have¹⁵ $\langle T_{ij} \rangle = N_i N_j / (N_T + \delta_{\beta 1})$, a result closely related to the Hauser-Feshbach formula in nuclear physics.²³ The WLC is then given by

$$\delta g = -N/N_\phi + O(1/N_\phi^2). \quad (7)$$

For the variance one finds from Eq. (5)

$$\text{varg} = \left(\frac{N}{N_\phi} \right)^2 \frac{2}{\beta} \left[1 + \frac{2-\beta}{2N} \right] + \dots \quad (8)$$

and, for the ratio of the variances for $\beta = 1, 2$,

$$\frac{(\text{varg})^{(\beta=1)}}{(\text{varg})^{(\beta=2)}} = 2 \left(1 + \frac{1}{2N} \right) + \dots \quad (9)$$

Of course the magnitude of the quantum corrections decreases as the number of phase-breaking channels increases. Note that the power controlling the decay of the variance is twice that of the average conductance. The deviation of the ratio of the variances from 2 is highly unusual; in fact, for $N = 1$ the ratio can be as high as 3.

We present in Figs. 1, 2 and 3 results of simulations of the random-matrix model in order to check the asymptotic behavior and suggest improvements. The points are obtained by generating random $N_T \times N_T$ unitary or orthogonal matrices and computing g from Eq. (1). In Fig. 1, the value $N = 2$ was selected for comparison with the experiment of Ref. 9. The log-log insets in Fig. 1 show that the convergence of the numerical to the asymptotic results [Eqs. (7)-(8), dotted lines] is rather slow. The curves in the main parts of Fig. 1 are interpolation formulae suggested by the results for $N_\phi = 0$ and $N_\phi \gg 1$. For the WLC, Eqs. (2a) and (7) suggest the relation

$$\delta g \simeq -N/(2N + N_\phi). \quad (10)$$

For the fluctuations of the conductance, we combine square roots of variances,

$$(\text{varg})^{1/2} = \left[(\text{varg})_{N_\phi=0}^{-1/2} + (\text{varg})_{N_\phi \gg 1}^{-1/2} \right]^{-1}, \quad (11)$$

because in the numerical simulations the change in varg is clearly linear in N_ϕ , not quadratic. So, for $\beta = 2$

$$\text{varg} \simeq N^2 / [(4N^2 - 1)^{1/2} + N_\phi]^2. \quad (12)$$

For $\beta = 1$ a similar, but more complicated, expression holds. These interpolation formulae agree very well with the numerical results; the only significant deviation is for $N = 1$ and small N_ϕ [Fig. 1(c)].

The results at fixed values of N_ϕ shown in Fig. 2 are relevant to experiments at fixed temperature in which the size of the opening to the cavity is varied. Though δg and varg are nearly independent of N in the perfectly coherent limit—a result known as “universality”—phase-breaking channels cause the magnitudes to vary. Thus the universality can only be seen if $N_\phi \ll N$; otherwise, the behavior is approximately linear, as in some

experiments.^{4,9} Clearly, phase-breaking must be included in interpreting the experiments.

In addition to the mean and variance, the probability density of the conductance, $w(g)$, is experimentally measurable.⁹ The numerical results in Fig. 3 show that as N_ϕ increases $w(g)$ tends towards a Gaussian and is therefore fully characterized by the mean and variance given above. For $N \geq 3$ the distribution is essentially Gaussian¹⁵ even for $N_\phi = 0$, so that N_ϕ affects $w(g)$, apart from changing the mean and variance, only for $N < 3$. For $N = 1$, the highly non-Gaussian distributions^{15,16} at $N_\phi = 0$ become approximately Gaussian at $N_\phi = 3$; the intermediate distributions, $N_\phi = 1, 2$, are shown in Fig. 3. For $N = 2$, the deviation of $w(g)$ for $N_\phi = 0$ from a Gaussian is smaller than for $N = 1$, and hence fewer phase-breaking channels produce a Gaussian distribution. We show numerical results only for $N_\phi = 1$.

Finally, we use our model to interpret the experiments of Refs. 4, 8 and 9. First, the solid circles in Fig. 1 show the measured N_ϕ , δg , and $\text{var} g$ of Ref. 9 in the $N = 2$ case. Before comparison the experimental variance must be corrected for thermal averaging: convolution over the derivative of the Fermi function produces a reduction factor of $\sim 0.22 - 0.38$ for an electron temperature of $50 - 100$ mK. We have increased the measured $\text{var} g$ by the inverse of this reduction factor; the WLC is not affected by thermal averaging since it is already an average effect. The error bars shown result from both the uncertainty in temperature in the case of the variance and the experimental fluctuations at small B for the variance and WLC. (For the moment, we do not assign further physical significance to the oscillations seen in the experiment.) Note that in Fig. 1 we have not fit the theory to the data and yet the agreement is very good. Second, in Fig. 2 we show as solid symbols the data of all three Refs. 4, 8 and 9. The value of N_ϕ deduced from comparing to our calculations is in good agreement with the value estimated independently in the experiments: $N_\phi = 7 - 9$ for Ref. 4, $N_\phi = 2$ for Ref. 8, and $N_\phi = 4 - 8$ for Ref. 9. Finally, the probability density of the conductance is found to be Gaussian in Ref. 9. For the estimated $N_\phi = 4 - 8$, we obtain a Gaussian distribution and so are consistent. Observation of the interesting non-Gaussian distribution of g obtained in the theory for $N = 1, 2$ and $N_\phi = 0$ requires greatly reduced phase-breaking.

In conclusion, we have presented a random-matrix model that simulates the effects of phase-breaking events on electronic transport through ballistic chaotic cavities. The analysis of recent experiments indicates that one can find a value of N_ϕ , the number of phase-breaking channels, that allows a consistent description of the data. Further experiments are needed to test quantitatively the dependence on N and N_ϕ that we predict.

We thank C. M. Marcus for several valuable conversations.

Note added— While preparing this paper for publication, we received a preprint by P. W. Brouwer and C. W. J. Beenakker with some overlapping material for $N = 1$.

-
- ¹ For a review see C. W. J. Beenakker and H. van Houten in *Solid State Physics*, Vol. 44, edited by H. Ehrenreich and D. Turnbull (Academic Press, New York, 1991) pp. 1-228.
 - ² C. M. Marcus, A. J. Rimberg, R. M. Westervelt, P. F. Hopkins and A. C. Gossard, *Phys. Rev. Lett.* **69**, 506 (1992).
 - ³ C. M. Marcus, R. M. Westervelt, P. F. Hopkins and A. C. Gossard, *Phys. Rev. B* **48**, 2460 (1993).
 - ⁴ M. J. Berry, J. H. Baskey, R. M. Westervelt and A. C. Gossard, *Phys. Rev. B* **50**, 8857 (1994); M. J. Berry, J. A. Katine, C. M. Marcus, R. M. Westervelt and A. C. Gossard, *Surf. Sci.* **305**, 480 (1994); M. J. Berry, *et al.*, preprint (Aug. 1994).
 - ⁵ M. W. Keller, O. Millo, A. Mittal, D. E. Prober, and R. N. Sacks, *Surf. Sci.* **305**, 501 (1994).
 - ⁶ D. Weiss, K. Richter, A. Menschig, R. Bergmann, H. Schweizer, K. von Klitzing, and G. Weimann, *Phys. Rev. Lett.*, **70**, 4118 (1993); D. Weiss, *et al.*, *Surf. Sci.* **305**, 408 (1994).
 - ⁷ R. Schuster, K. Ensslin, D. Wharam, S. Kühn, J. P. Kotthaus, G. Böhm, W. Klein, G. Tränkle, and G. Weiman, *Phys. Rev. B* **49**, 8510 (1994).
 - ⁸ A. M. Chang, H. U. Baranger, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **73**, 2111 (1994).
 - ⁹ I. H. Chan, R. M. Clarke, C. M. Marcus, K. Campman, and A. C. Gossard, preprint (Aug. 1994); R. M. Clarke, I. H. Chan, C. M. Marcus, C. I. Duruöz, J. S. Harris Jr., K. Campman, and A. C. Gossard, preprint (Aug. 1994).
 - ¹⁰ R. Blümel and U. Smilansky, *Phys. Rev. Lett.* **60**, 477 (1988); **64**, 241 (1990); *Physica D* **36**, 111 (1989).
 - ¹¹ R. A. Jalabert, H. U. Baranger, and A. D. Stone, *Phys. Rev. Lett.* **65**, 2442 (1990); H. U. Baranger, R. A. Jalabert, and A. D. Stone, *Phys. Rev. Lett.* **70**, 3876 (1993); *Chaos* **3**, 665 (1993).
 - ¹² A. Pluhar, H. A. Weidenmüller, and J. A. Zuk, *Phys. Rev. Lett.* **73**, 2115 (1994).
 - ¹³ K. Efetov, preprint (July 1994).
 - ¹⁴ S. Iida, H. A. Weidenmüller, and J. A. Zuk, *Phys. Rev. Lett.* **64**, 583 (1990) and *Annals of Phys.* **200**, 219 (1990).
 - ¹⁵ H. U. Baranger and P. A. Mello, *Phys. Rev. Lett.* **73**, 142 (1994).
 - ¹⁶ R. A. Jalabert, J.-L. Pichard, and C. W. J. Beenakker, *Europhys. Lett.* **27**, 255 (1994); R. A. Jalabert and J.-L. Pichard, preprint (September 1994).
 - ¹⁷ M. Büttiker, *Phys. Rev. B* **33**, 3020 (1986).
 - ¹⁸ B. Schmidt and A. Müller-Groeling, *Phys. Rev. B* **47**, 12732 (1993) and references therein.
 - ¹⁹ E. Doron, U. Smilansky, and A. Frenkel, *Physica D* **50**, 367 (1991).
 - ²⁰ For a review see M. L. Mehta, *Random Matrices* (Academic, New York, 1991).
 - ²¹ P. Pereyra and P. A. Mello, *J. Phys. A* **16**, 237 (1983).
 - ²² P. A. Mello, *J. Phys. A* **23**, 4061 (1990).
 - ²³ W. A. Friedman and P. A. Mello, *Ann. Phys. (N.Y.)* **161**, 276 (1985) and references therein.

FIG. 1. Magnitude of quantum transport effects as a function of the number of phase-breaking channels, N_ϕ , on linear (main panels) and log-log (insets) scales. (a) The weak-localization correction ($N = 2$). (b) The variance in both the orthogonal (squares) and unitary (triangles) cases ($N = 2$). (c) The ratio of the variance in the orthogonal case to that in the unitary for $N = 1, 2$, and 6; the arrows mark the $N_\phi \rightarrow \infty$ limit for $N = 1, 2$. Open symbols are numerical results (20,000 matrices used, statistical error the size of the symbol). Solid lines are interpolation formulae. Dotted lines are asymptotic results. Solid circles are experimental results of Ref. 9 corrected for thermal averaging. The interpolation formulae are excellent except for $N = 1$ and small N_ϕ [panel (c)].

FIG. 2. The magnitude of quantum transport effects as a function of the number of channels in the leads, N , for fixed $N_\phi = 0, 2, 4$, and 8. (a) The weak-localization correction. (b) The variance for the orthogonal case ($B = 0$). (c) The variance for the unitary case (nonzero B). Open symbols are numerical results (as in Fig. 1). Solid lines are interpolation formulae. Solid symbols are experimental results of Refs. 4 (triangles), 8 (squares), and 9 (circles) corrected for thermal averaging. The introduction of phase-breaking decreases the “universality” of the results but leads to good agreement with experiment.

FIG. 3. The probability density of the conductance in the orthogonal (first column) and unitary (second column) cases for $N = 1$ (first row) and $N = 2$ (second row). Increasing the phase-breaking from zero (dashed lines, analytic) to $N_\phi = 1$ (plus symbols, numerical) to $N_\phi = 2$ (squares, numerical) moves the distribution towards a Gaussian.




Figure 1.




Figure 2.




Figure 3.